

LETTER • OPEN ACCESS

Managing forest carbon and landscape capacities

To cite this article: Céline Boisvenue *et al* 2022 *Environ. Res. Lett.* **17** 114013

View the [article online](#) for updates and enhancements.

You may also like

- [Development and investigation of densification behavior of ZrB₂-SiC composites through microwave sintering](#)
Ankur Sharma and D B Karunakar
- [Limits to forests-based mitigation in integrated assessment modelling: global potentials and impacts under constraining factors](#)
Théo Rouhette, Neus Escobar, Xin Zhao et al.
- [Prospective regularization design in prior-image-based reconstruction](#)
Hao Dang, Jeffrey H Siewerdsen and J Webster Stayman

UNITED THROUGH SCIENCE & TECHNOLOGY



The Electrochemical Society
Advancing solid state & electrochemical science & technology

248th ECS Meeting

Chicago, IL
October 12-16, 2025
Hilton Chicago



**Science +
Technology +
YOU!**

Abstract submission
deadline extended:
April 11, 2025

SUBMIT NOW

ENVIRONMENTAL RESEARCH
LETTERS

LETTER

Managing forest carbon and landscape capacities

Céline Boisvenue^{1,2,4,*} , Gregory Paradis² , Ian M S Eddy¹ , Eliot J B McIntire^{1,2,4}
and Alex M Chubaty^{3,4} ¹ Pacific Forestry Centre, Canadian Forest Service, Natural Resources Canada, 506 Burnside Rd W, Victoria, BC V8Z 1M5, Canada² Faculty of Forestry, Forest Resources Management, The University of British Columbia, 2045-2424 Main Mall, Vancouver, BC V6T 1Z4, Canada³ FOR-CAST Research & Analytics, PO BOX 96026 West Springs, Calgary, AB T3H 0L3, Canada⁴ Current address/primary affiliation.

* Author to whom any correspondence should be addressed.

E-mail: celine.boisvenue@nrcan-rncan.gc.ca**Keywords:** nature-based mitigation, forest carbon, forest management, carbon carrying -capacitySupplementary material for this article is available [online](#)

RECEIVED

22 November 2021

REVISED

13 September 2022

ACCEPTED FOR PUBLICATION

11 October 2022

PUBLISHED

21 October 2022

Original content from
this work may be used
under the terms of the
[Creative Commons
Attribution 4.0 licence](#).

Any further distribution
of this work must
maintain attribution to
the author(s) and the title
of the work, journal
citation and DOI.



Abstract

Widespread impacts of a warming planet are fuelling climate change mitigation efforts world-wide. Decision makers are turning to forests, the largest terrestrial primary producer, as a nature-based contribution to mitigation efforts. Resource-based economies, however, have yet to include carbon (C) in their resource planning, slowing the implementation of these important measures for atmospheric greenhouse gas reduction. The realisation of forest mitigation potential depends greatly on our ability to integrate C-sequestration practices in our forest management applications. This requires robust C-estimates, an understanding of the natural potential for a specific landscape to sequester C, the current state of the landscape relative to this potential, and the evaluation of management practices as a tool to sequester forest C in the midst of all the other values forests offer humans. Discrepancies between models used in management decisions and C estimation are the first hurdle impeding the application of forest-based mitigation strategies. Here, we combine forest disturbance and management models with a well-established C model on an open-source simulation platform. We then use the modelling system to produce C estimates of the natural C-holding capacity (potential) and two management scenarios for a study area in BC, Canada. Our simulations provide an essential metric if forests are to be managed for C-sequestration: the natural landscape C-holding capacity. Our simulations also point to a decreasing trend in simulated C on the study area over time and to a bias of the current C-levels compared to the landscape C-holding capacity (477 vs 405.5 MtC). Our explanations for this bias may provide an avenue for improved current C-state estimates. We provide a framework and the information needed for the implementation of nature-based solutions using forests for climate change mitigation. This study is a step towards modelling systems that can unify scientifically based forest management and informed C-management.

1. Introduction

Forests are the largest terrestrial primary producer, absorbing the largest proportion of atmospheric CO₂ (Friedlingstein *et al* 2020). They have considerable potential to help mitigate the impacts of changing climatic conditions (Griscom *et al* 2017, Cook-Patton *et al* 2020). To realise this potential however, C needs to be an integral part of forest management decisions. Impediments to including C in forest management

decisions vary per jurisdiction, but have a common trait: discrepancies between models used in management decisions and C estimation. The sheer complexity of the forest C cycle makes it challenging for measuring, monitoring, and projecting changes to forest C (Boisvenue and White 2019, Forests in focus 2021). Further, the geographic extent and spatiotemporal variability in the system prohibits, in most cases, a complete census of forest C (FAO 2019), which implies that sampling and modelling are therefore

essential to forest management and planning. Projecting productivity via modelling is the basis for sustainable forest management and modelling of species dynamics supports silvicultural interventions. Modelling efforts have also resulted in forest C and its variations being estimated with a diversity of models (Xia *et al* 2017, Walker *et al* 2022). Despite these efforts, gaps remain in our theoretical and empirical understanding of C allocation (Merganičová *et al* 2019).

The call for integration of forest-based solutions in emission reduction policy is increasing (Anderegg *et al* 2020, Matthews *et al* 2020, Lamb *et al* 2021) with policies being put in place for integration of forest-based intervention in the mitigation efforts (Environment and Climate Change Canada 2020). Their application, where management actions aimed at increased C storage are implemented, are still restricted to academic publications (Liu *et al* 2016, Pilli *et al* 2021, Sleeter *et al* 2022, Zhao *et al* 2022)⁵. Even in jurisdictions where forest C is quantified, most do not include forest C when planning greenhouse gas reductions (Lamb *et al* 2021), nor is C included in forest management decisions⁶. This is partly because our ability to manipulate the C content of forests rests within the realm of forest management, and models supporting those decisions differ in structure and spatial extent from forest C models (e.g. Fortin and Lavoie 2022 vs Walker *et al* 2022). Nonetheless, forest-based interventions need C information and decision makers face these modelling discrepancies amongst other challenges (Grassi *et al* 2017, Markusson 2022). Sequestration potentials have been estimated using C-models (Griscom *et al* 2017, 2020, Smyth *et al* 2020, Drever *et al* 2021). In Canada for example, forest and specifically improved forest management is estimated to offer the 4th largest mitigation opportunities for 2030 among natural climate solutions (Drever *et al* 2021). Even with a well-established forest C model, as is the case for Canada (Environment and Climate Change Canada 2020), forest C management is, as of yet, not incorporated into resource planning. Effective forest-based solutions require (a) robust estimates of current C, (b) an assessment of the land's potential and unrealised potential for C-sequestration, and (c) integration of C estimates and potential into land-management decision schema (Walker *et al* 2022).

Landscape characteristics partly determine the amount of C that can be maintained, i.e. its potential. This amount varies through time, and is bounded by environmental and site conditions (Boisvenue and Running 2006, Running 2012, Walker *et al* 2022).

The potential or C-holding capacity is the mass of C stored in a landscape under prevailing historical environmental conditions and natural disturbance regimes (Keith *et al* 2010, Liang *et al* 2017) excluding anthropogenic interventions. A disturbance regime is a historic pattern (e.g. frequency and extent) of natural processes such as fire, insects, wind, and mass movement that affect the ecosystems and landscapes in a particular area. The C-holding capacity is a theoretical value that reflects the long-term stabilization of the landscape C under the dominant natural disturbance. If the dominant disturbance is fire, this C-potential is linked to the historic fire return interval (FRI). Current potential to hold C is a legacy of past environmental conditions. It will change with changing environmental conditions as those drive the disturbance regimes. Most forest-based economies have legislated sustainable forestry practices⁷, but planning horizons are short, varying from five to ten years⁸. Given the short forest management time horizon, the notion of current capacity is necessary for informed C-management decisions. Longer-term planning will need to be informed by projections and probabilities of changing disturbance regimes under changing environmental conditions, a field of research not yet ripe for integration in shorter-term forest management decisions. Many C-centric studies compare various management approaches but omit the notion of current C-holding capacity (Pilli *et al* 2016, Krug 2019). Other studies assess the potential of the landscape to hold C but do not link it to the decision-making process involved in forest resource management (Keith *et al* 2010, Wei *et al* 2014, Domke *et al* 2020).

Given current political and environmental motivations to maximize our climate change mitigation capacities, quantitative assessments of C in the landscape, of management interventions C-effects, and of landscape C-holding capabilities are in urgent need. Here, we develop a modelling system, combining forest disturbance and management simulations with a well-established C model for a large study area in western Canada. We then apply this system to (a) estimate C, and (b) calculate the current C-holding potential of a landscape. We then compare current potential to current conditions, and two scenarios of resource removal and discuss our findings. We follow modelling best practices of McIntire *et al* (2022) resulting in our approach being transparent, repeatable and reusable. We provide an avenue for implementing forest-based greenhouse gas reduction

⁵ No C in resource values: www2.gov.bc.ca/gov/content/industry/forestry/managing-our-forest-resources/forest-stewardship-plans.

⁶ Example: www2.gov.bc.ca/gov/content/industry/forestry/managing-our-forest-resources/timber-supply-review-and-allowable-annual-cut.

⁷ Example: www.nrcan.gc.ca/our-natural-resources/forests/sustainable-forest-management/canadas-forest-laws/17497.

⁸ Example: www2.gov.bc.ca/gov/content/industry/forestry/forest-tenures/forest-tenure-administration/apportionment-commitment-reports-aac#:~:text=The%20allowable%20annual%20cut%20commonly,least%20once%20every%20ten%20years.

interventions, and transparent tools to evaluate and eventually improve forest C management.

2. Methods

2.1. Study area

We selected a study area in British Columbia, Canada, a forest-based economy⁹ where a reputable approach is used for forest C estimation and reporting (Kurz *et al* 2009, Stinson *et al* 2011, Metsaranta *et al* 2017)¹⁰. Our study area is located at the northeastern corner of the province of British Columbia (figure 1). It encompasses five forest management units in the provincial forest management system with 31 Mha of 53.4 Mha region under provincial forest management agreements. The topography of the units forms a gradient of increasing relief from east to west. Under provincial legislation (Forest Act 1996¹¹), these lands are actively managed for the sustainable production of resources including wood products, with a combined 19 Mm³ yr⁻¹ of annual allowable harvest¹².

2.2. Modelling

2.2.1. Context

Even in places like Canada, where the Federal Government reports annually on greenhouse gas emissions under the United Nations Framework Convention on Climate Change¹³ for all managed forests, C modelling is not linked to forest management modelling. In this case, C modelling uses similar growth information as forest management models, however, incompatibilities in the software and information processing exist, and a mismatch in spatial resolution between the modelling systems results in C information being currently external to the modelling used in forest resource management. Growth modelling has emerged from the history of resource management for wood supply. Modelling of forest C was developed for the international greenhouse gas reporting, with Canada submitting its first report 2006¹⁰. In our example, the Chief Forester of BC determines annual allowable cut (AAC) level¹⁴, informed by growth and yield modelling which represents forest dynamics. In the case of our study area, these models are the basis for the C-estimates produced by the Canadian Forest Service, who is responsible for reporting on the

greenhouse gases (GHG) balance. For GHG reporting, growth models inputs and outputs are processed differently and applied at a coarser scale than that needed on forest management decisions (Kurz *et al* 2009). Here we start with the model currently used for reporting, the Carbon Budget Model of the Canadian Forest Sector (CBM—Kurz *et al* 2009), adapting it so it can provide C-estimates compatible with forest management decisions. We then link the C-modelling with a fire model to enable the calculation of the C-holding capacity of the landscape, since fires are the dominant natural disturbances in our study area (Kurz *et al* 2013). Finally, we add a forest extraction model to assess scenarios of resource extraction and their effects on C.

Our modelling efforts strive towards an open reusable approach, as described in McIntire *et al* (2022). All three models were re-implemented or linked via *SpaDES* (**S**patial **D**iscrete **E**vent **S**imulation—McIntire *et al* 2019), an open source platform that enables model connections and expansion to a spatially explicit format. This platform offers a solid framework to build interoperable and reusable models (e.g. Micheletti *et al* 2021) in the R language, a free software environment¹⁵. The source code used for this study is publicly available on GitHub¹⁶ and follows FAIR data standards (Findable, Accessible, Interoperable and Reusable data; Stall *et al* 2019).

2.2.2. Forest carbon modelling

An executable for CBM is freely available¹⁷, but runs on Microsoft Access, which in many cases limits a user's ability to run, modify and understand the model and data processing. We therefore re-implemented CBM in R, on the *SpaDES* platform (Supplementary information Appendix 1). The implementation of CBM in R and *SpaDES* enabled modifications to the spatial resolution of simulations, making it more flexible and able to match a spatial resolution of forest management models. It also renders the model and data processing transparent, enabling us to address the incompatibilities in the software and information processing between the C model and the forest management models. CBM has parameters for all managed forests of Canada (Stinson *et al* 2011), which we use for our study area and make available for all of Canada in our modelling system. CBM derives live aboveground biomass estimates, which drive its growth and post-disturbance dynamics, using field-measured yield information from the forest industry (m³ ha⁻¹ over time) translated into biomass using statistical equations (Boudewyn *et al* 2007). Other C pools in CBM are estimated via process

⁹ www.nrcan.gc.ca/our-natural-resources/forests/state-canadas-forests-report/forest-industry-contribute/16517.

¹⁰ www.nrcan.gc.ca/climate-change-adapting-impacts-and-reducing-emissions/climate-change-impacts-forests/forest-carbon/13085.

¹¹ www.bclaws.gov.bc.ca/civix/document/id/complete/statreg/96157_00

¹² www2.gov.bc.ca/gov/content/industry/forestry/managing-our-forest-resources/timber-supply-review-and-allowable-annual-cut/allowable-annual-cut-timber-supply-areas.

¹³ unfccc.int/process-and-meetings/transparency-and-reporting/greenhouse-gas-data/ghg-data-unfccc/ghg-data-from-unfccc.

¹⁴ Example: Fort Nelson Timber Supply Area news.gov.bc.ca/releases/2019FLNR0187-001442.

¹⁵ www.r-project.org/.

¹⁶ github.com/cboisvenue/spadesCBM_RIA.git.

¹⁷ www.nrcan.gc.ca/climate-change/impacts-adaptations/climate-change-impacts-forests/carbon-accounting/carbon-budget-model/13107.




Figure 1. Our study area, in British Columbia, Canada.

representation during an initialisation phase. The initialisation phase aims at stabilizing C in dead organic matter pools. Parameters and processes for CBM are described in detail in Kurz *et al* (2009). CBM tracks C in 21 pools for each unit (stand or pixel), plus three pools for emissions (CO_2 , CH_4 , CO), and one for C removed from the system via harvested wood products. The flexibility and transparency enabled by R and *SpaDES* (McIntire *et al* 2019, 2022) permitted a direct link to the management and fire simulator resulting in the management and C models being driven by the same growth models, sharing input data and data processing.

2.2.3. Resource extraction modelling

Two management scenarios were simulated using a *SpaDES* adaptation of the *ws3* model, a Python-based open source software package (Paradis and LeBel 2018). *Ws3* reproduces industry-standard forest estate modelling functionality, used by foresters to determine AAC. We used *ws3* to generate plausible patterns of harvesting and regeneration for the managed portions of each timber supply area (TSA). The *spades_ws3* module (Supplementary information Appendix 2) wraps *ws3* in R scripts to yield a *SpaDES*-compatible module that we used in our study to combine with CBM.

2.2.4. Fire modelling

We used a *SpaDES* implementation of the fire model described in Armstrong and Cumming (2003) (Supplementary Information Appendix 3). This model treats fire as a three-stage stochastic process of ignition, escape, and spread adapted to the specific landscape characteristics. To eliminate discrepancies between the fire simulation and the C-estimation, these processes had to be parameterized to simulate the FRI used in the CBM initialization phase. Disturbance-return intervals used for CBM initialization differ for each terrestrial ecozone and vary between 125 and 175 years in our study area (see Kurz *et al* 2009).

2.3. Data

We maintained FAIR data standards (Stall *et al* 2019) as much as possible for our input data and used publicly available parameters for our simulations. Our data preparation modules¹⁸, which we developed for each scenario, provide links to all publicly available data.

We used the default Canadian parameters for CBM (Kurz *et al* 2009, Stinson *et al* 2011) that

¹⁸ github.com/cboisvenue/spadesCBM_RIA/tree/main/modules.

are specific to our study area. British Columbia has an open access policy and we used their Vegetation Resource Inventory to populate our forest stand ages for each of the scenarios and across models. All forest inventory stands in CBM are represented by the growth curve of the dominant species, reflecting the predominantly stand-replacing impacts of boreal disturbance regimes. Stands can contain hardwood and softwood components, each associated with individual growth information. The growth curves used for forest management are not publicly available. We simplified the forest management growth curves (described in Supplementary Information Appendix 2) so we could provide all inputs to our system, making this modelling system and simulations transparent. We provide a link in our data preparation module to the simplified curves¹⁴. Disturbance information was specific to each scenario and is described in the following section.

2.4. Simulations scenarios

2.4.1. Landscape C-holding capacity

Disturbed stands can be expected to be emitters of atmosphere greenhouse gases either because of combustion or because of decay of organic material or both. Under a disturbance regime, a landscape can maintain a relatively stable amount of C (Keith *et al* 2010, Liang *et al* 2017). To estimate this C-holding capacity for our study area, we simulated fire at return intervals that matched the initializing phase used in CBM, with no other disturbances, starting from the British Columbia Vegetation Resource Inventory ages and dominant species in 2020, for 520 years. This accounted for the stochasticity in fire ignition, escape and spread, as well as the stabilization of the C on the landscape. The location of these fire disturbances were the product of our fire model (see section 2.3) and account for the characteristics of our study area. The representation of the C pool impacts of the disturbances in each scenario are those described in CBM and specific to our location in Canada (see Kurz *et al* 2009, Stinson *et al* 2011).

2.4.2. Present day

We simulated a representation of present-day conditions for our study area by simulating the same landscape, using the same modelling framework but for the time-period 1985–2015. By simulating a present-day scenario, we have a representation of current conditions through the lens of our modelling system which we use as a comparison point. Growth information for this scenario matched that used in other scenarios. Landscape conditions were derived from the BC Vegetation Resource Inventory with stand replacing disturbances locations from National Forest Information System¹⁹ (Hermosilla *et al* 2016). As per all C-implications of disturbances in our scenarios,

disturbances followed their specific C transactions prescribed in CBM for this location in Canada (see Kurz *et al* 2009, Stinson *et al* 2011).

2.4.3. Harvesting scenarios

We simulated two resource removal scenarios one higher ('base') and one lower ('less') from the year 2020–2099. This time horizon represents multiple planning cycles in forest management⁸. The *base* scenario simulates harvesting the maximum sustainable even-flow volume at every time step. As a comparison point, and to corroborate the general pattern of increased resource extraction corresponding to a decrease in C storage from the scientific literature (Harmon and Marks 2002, Bradford 2011, Sharma *et al* 2013, Simard *et al* 2020), the *less* scenario simulates harvesting half of the volume from the *base* scenario. Both scenarios extract resources from the landscape via whole-stand disturbance over the horizon period as estimated by *spades_ws3*. *Spades_ws3* schedules these events along with an expected level of wild-fires, supplying the location of each stand-replacing disturbance to the modelling system. Under the *less* scenario, we expected that more C would be maintained on the landscape.

3. Results

Our simulations estimate a C-holding capacity of 405.5 MtC for our study area (table 1, figure 3 panel (a)). While all scenarios showed a decrease in total carbon from the start to the end of simulations (Table 1), they all started with landscape-level C above the C-holding capacity (table 1). The distribution of C in the landscape differs between scenarios with the present-day and harvest scenarios having fewer pixels with low tC/ha but more with mid-range tC/ha than the C-holding capacity scenario (figure 2). The present-day scenario ended simulations with landscape-level C higher than C-holding capacity (477.2 MtC), while the two harvest scenarios ended with C lower than C-holding capacity. Harvesting less retained more carbon on the landscape (Δ 33.7 MtC) and although estimates from both harvesting scenarios kept less C on the landscape than the estimated C-holding capacity, the lesser-level of resource removal was close to C-holding capacity.

The average influx of C per ha under the C-holding capacity scenario stabilises around the 480th year of simulation at ~ 0.60 MtC ha yr⁻¹ (table 2). The *less* harvest scenario C uptake has the same pattern as the present-day simulations, with a decreasing level of C-uptake through the simulation horizon. The *base* harvest case maintains a higher net C-uptake. Current levels of resource removals as represented in our simulations, are lower than the representation of maximum sustainable removal of wood products (*base*). When only looking at the

¹⁹ opendata.nfis.org/mapserver/nfis-change_eng.html.

Table 1. Total carbon, above and belowground, in MtC for each scenario at the start of simulations (start = landscape initialization results) and at the end.

Scenario	Simulation	
	Start	End
C-holding capacity	560.5	405.5 ^a
Present day	556.3	477.2
Base harvest	560.5	367.3
Less harvest	560.5	400.9

^a C-holding capacity.




Figure 2. Density plots of the tC per ha in the pixels for each of the scenarios at the end of the simulations: C-holding capacity, present day, base harvest and less harvest.

Table 2. Net primary productivity in MtC ha⁻¹ for each scenario at the start and end of each simulation.

Scenario	Simulation	
	Start	End
C-holding capacity	0.723	0.60
Present day	0.718	0.63
Base harvest	0.718	0.70
Less harvest	0.723	0.64


C in aboveground biomass, the change over the C-holding capacity simulation is very close to the change in the *less harvest* scenario (116 MtC and 113 MtC respectively).

4. Discussion

Our modelling system linked C, disturbance and management models to produce C estimates, C-holding capacity, and two levels of resource


removal in our study area (figure 3). C-holding capacity simulations display a negative exponential age class distribution (figure 4(a); Wagner 1978), indicating that our goal of stabilizing landscape-C was achieved. How much C a landscape can naturally hold is an essential metric if forests are to be managed for C-sequestration (Griscom *et al* 2017, 2020, Drever *et al* 2021, Walker *et al* 2022).

Our simulations show a decreasing trend in C-retention, with all initialized as well as the present-day (2015) landscapes having higher C than the C-holding capacity. Given the conceptual similarities between the C-holding capacity simulations and the initialization process (i.e. both repeat a grow-burn cycle), their results should be similar. Fire suppression or management are not accounted for in CBM initialization process further indicating that both should yield similar results. They differ, however, in that the CBM-initialization process burns stands at regular intervals equal to the FRI and our C-holding



simulations burn at stochastic intervals with a mean return interval of FRI. This results in an upwards biased initial C estimate because of the mathematical reality of asymmetric C accumulation in growth curves on either side of the FRI age. In other words, if a stochastic fire burns 80 years younger than FRI, the stand will have *not* accumulated X amount of C, while the amount of C gained due to 80 years of extra growth beyond FRI age is $a \times X$, where $a < 1$. This effect seems to be insensitive to the particular distribution of FRI used as it will occur as long as the growth is slowing at FRI age (i.e. negative exponential or normally distributed around a mean of FRI; see Supplementary Information Appendix 4). This lower C storage (greater loss) due to more burns younger than FRI age has recently been noticed in nearby northern Canadian forests (Baltzer *et al* 2021). In our C-holding capacity simulations, burns will

stochastically occur during the pre-FRI age, resulting in a C accumulation that is cut short compared to the FRI age. Because the growth curves prescribe slowing of growth past FRI age, the stochastically longer-than-FRI-age inter-burn periods do not make up for the many instances of pre-FRI burns (see Supplementary Information Appendix 4). We also see a narrower age distribution (figures 4(a) and (c)), which is likely either the result of these forests being outside of their natural range due to forest management that uses a fixed rotation age or inaccurate age estimates especially of the older trees (Cumming *et al* 2000). This also may contribute to why initialized landscapes have higher C than the C-holding capacity. The initialization step has been necessary because we have lacked data on the highly variable soil and dead C-pools. Varying the burn interval in the CBM-initialization process has been shown to be a source of variation



(Metsaranta *et al* 2017) and initialization processes in other contexts have been shown to be the source of systematic errors (Oberpriller *et al* 2021). With the increase in availability of soils and belowground C pools data (Sothe *et al* 2022), initialization processes may eventually become unnecessary. The comparison of the landscape-state to its C-holding capacity will not only require more exploration of this initialization process, but will probably necessitate data on C-content of dead organic layers.

In our simulations, both harvesting scenarios have the same initial landscape-C values. Even if these were wrong, the comparison of resulting

landscape-C would still be valid. Our results show that the higher extraction in our base-scenario results in less landscape-C than the lower extraction level (Δ 33.6 MtC) matching other published trends (Harmon and Marks 2002, Bradford 2011, Sharma *et al* 2013, Simard *et al* 2020). Some mitigation scenarios suggest storing C outside the system, such as in durable harvested wood products, and keeping the system at a higher influx (higher C-influx in the *base* scenario table 2, Smyth *et al* 2020). Assessing this scenario would require adding a wood product and life cycle analysis model to our modelling system, an opportunity presented by our modular and flexible system. C-storage outside the system may justify a higher harvest level through C-sequestration in wood products, but may have other undesired consequences such as lowering landscape resilience (Ibáñez *et al* 2019, Albrich *et al* 2020) and soil C impacts (Prescott *et al* 2021). Further, higher influx into the system does not necessarily equal C-residency (Körner *et al* 2005).

Both harvesting scenarios result in landscape-C below C-holding capacity, although the *less* harvest scenario is closer to the C-holding capacity (Δ <5 MtC over the 53.4 Mha). Assuming that our landscape C-holding capacity approximates the actual holding capacity and that our harvesting simulations represent the correct trends, this simulation study suggests that, if only landscape-C is considered, extracting a lower amount of resource may match the landscape-C to its C-holding capacity. However, we have not yet determined the consequences of maintaining a landscape above or below its natural C-holding capacity (for example, for an increased C-storage elsewhere). Our current landscape (2015) has higher C than the C-holding capacity. This could suggest that the landscape is holding more C than what it could under the assumed-constant natural disturbance regime, potentially due to its historical anthropogenic interventions. It could also be stabilizing after too-high C-values from the initialization phase (see above). Given the uncertainty about the initialization process, to really determine if a landscape can hold more than its C-holding capacity and the consequences of maintaining a landscape in that state, more research is required. Our system is well-positioned to advance this research.

Our simulations indicate that the stable-state of the forests may be better captured by improved fire modelling. A stable-state, however, may be an irrelevant concept under changing environmental conditions which would require forecasts of evolving disturbance regimes in our landscapes (Walter *et al* 2019). Further, most forested regions are not pristine areas devoid of human influence, but are places where people have played a role in shaping the trajectory of forest composition, and thus C storage, over time (Lewis *et al* 2015), be it only via fire suppression (Hurteau *et al* 2019).

5. Summary and concluding remarks

We successfully adapted CBM to match the spatial scale of the management model, combined CBM with a fire model that uses the characteristics of the landscape to simulate the current disturbance regime until the landscape-level C stabilized, and standardized the software and information processing across all three models. This modelling system attempts to follow the reproducible, transparent and reusable approach as per McIntire *et al* (2022) rendering it readily adaptable to other study areas and available to iterative updating (Dietze *et al* 2018). Given that forest management and C models are jurisdiction-specific, our example is most easily reusable in BC, can be adapted to the rest of Canada, but could be adapted to jurisdictions with only the jurisdiction-specific information modified, rather than the whole workflow. At a minimum, it provides a transparent blueprint to linking models for C-inclusion in management decisions.

The realisation of forest mitigation potential depends greatly on our ability to integrate C-sequestration practices in our forest management applications. This requires robust C-estimates, an understanding of the potential for a specific landscape to sequester C, the current landscape-state relative to this potential, and the evaluation of management practices as C-sequestration tools in the midst of all the other commodities forests offer humans. Modelling efforts must strive to disentangle forest responses. Our study presents a solid basis for such an effort. Improved representation of our systems' current state will require improved data sets, the ability to iteratively update and improve simulations (as per McIntire *et al* 2022), which may even enable projections of the potential states of forests (Trouillier *et al* 2020) while using models that can support applied decisions. Our study is a step towards modelling systems that can support scientifically based decision-making and informed C-management.

Data availability statement

The data that support the findings of this study are openly available at the following URL/DOI: https://github.com/cboisvenue/spadesCBM_RIA.git.

Funding

This research was undertaken as a project in the *Science-based improvements to the National Forest Carbon Monitoring, Accounting and Reporting System*, in the work package *Forest carbon tools and assessments for mitigation and reporting*, in the *Forest Climate Change* program of the Canadian Forest Service.

Statement of authorship

C B conceived of the idea, completed the joint simulations, and wrote the manuscript; E M and C B completed the analyses and jointly developed the discussion; E M and A C are lead developers of the *SpaDES* toolkit, tested and helped adapt models to the modelling system; G P developed the harvesting module (*ws3*), participated in the conception of the research, provided harvest scenario simulations and growth representation; I E implemented and provided the C-holding capacity fire simulations and provided GIS support; all authors contributed substantially to revisions.

ORCID iDs

Céline Boisvenue  <https://orcid.org/0000-0002-6031-7961>

Gregory Paradis  <https://orcid.org/0000-0001-9618-8797>

Ian M S Eddy  <https://orcid.org/0000-0001-7397-2116>

Eliot J B McIntire  <https://orcid.org/0000-0002-6914-8316>

Alex M Chubaty  <https://orcid.org/0000-0001-7146-8135>

References

- Albrich K, Rammer W, Turner M G, Ratajczak Z, Braziunas K H, Hansen W D and Seidl R 2020 Simulating forest resilience: a review *Glob. Ecol. Biogeogr.* **29** 2082–96
- Anderegg W R L *et al* 2020 Climate-driven risks to the climate mitigation potential of forests *Science* **368** eaaz7005
- Armstrong G W and Cumming S G 2003 Estimating the cost of land base changes due to wildfire using shadow prices *For. Sci.* **49** 719–30
- Baltzer J L *et al* 2021 Increasing fire and the decline of fire adapted black spruce in the boreal forest *Proc. Natl Acad. Sci.* **118** e2024872118
- Boisvenue C and Running S W 2006 Impacts of climate change on natural forest productivity—evidence since the middle of the 20th century *Glob. Change Biol.* **12** 862–82
- Boisvenue C and White J C 2019 Information needs of next-generation forest carbon models: opportunities for remote sensing science *Remote Sens.* **11** 463
- Boudewyn P, Song X, Magnussen S and Gillis M D 2007 Model-based, volume-to-biomass conversion for forested and vegetated land in Canada (P. F. C. Canadian Forest Service, Trans; Information Report BC-X-411). Natural Resources Canada
- Bradford J B 2011 Potential influence of forest management on regional carbon stocks: an assessment of alternative scenarios in the Northern Lake States, USA *For. Sci.* **57** 479–88
- Cook-Patton S C *et al* 2020 Mapping carbon accumulation potential from global natural forest regrowth *Nature* **585** 545–50
- Cumming S G, Schmiegelow F K A and Burton P J 2000 Gap dynamics in boreal aspen stands: is the forest older than we think? *Ecol. Appl.* **10** 744–59
- Dietze M C *et al* 2018 Iterative near-term ecological forecasting: needs, opportunities, and challenges *Proc. Natl. Acad. Sci.* vol 115 pp 1424–32

- Domke G M, Oswalt S N, Walters B F and Morin R S 2020 Tree planting has the potential to increase carbon sequestration capacity of forests in the United States *Proc. Natl Acad. Sci.* **117** 24649–51
- Drever C R *et al* 2021 Natural climate solutions for Canada *Sci. Adv.* **7** eabd6034
- Environment and Climate Change Canada 2020 Climate science 2050: advancing science and knowledge on climate change Monograph No. En4-414/2020E-PDF 62 (Government of Canada) (available at: <http://publications.gc.ca/pub?id=9.892783&sl=0>)
- FAO 2019 State of the world's forests (forests and agriculture: land-use challenges and opportunities) (available at: www.fao.org/publications/sofo/2016/en/)
- Forest Act 1996 (available at: https://www.bclaws.gov.bc.ca/civix/document/id/complete/statreg/96157_00)
- Forests in Focus 2021 *Nat. Clim. Change* **11** 363
- Fortin M and Lavoie J-F 2022 Reconciling individual-based forest growth models with landscape-level studies through a meta-modelling approach *Can. J. For. Res.* **52** 1140–53
- Friedlingstein P *et al* 2020 Global carbon budget 2020 *Earth Syst. Sci. Data* **12** 3269–340
- Grassi G, House J, Dentener F, Federici S, den Elzen M and Penman J 2017 The key role of forests in meeting climate targets requires science for credible mitigation *Nat. Clim. Change* **7** 220–6
- Griscom B W *et al* 2017 Natural climate solutions *Proc. Natl Acad. Sci.* **114** 11645
- Griscom B W *et al* 2020 National mitigation potential from natural climate solutions in the tropics *Phil. Trans. R. Soc. B* **375** 20190126
- Harmon M E and Marks B 2002 Effects of silviculture practices on carbon storage in Douglas-fir-western hemlock forest in the Pacific Northwest, U.S.A.: results from a simulation model *Can. J. For. Res.* **32** 863–77
- Hermosilla T, Wulder M A, White J C, Coops N C, Hobart G W and Campbell L B 2016 Mass data processing of time series Landsat imagery: pixels to data products for forest monitoring *Int. J. Digit. Earth* **9** 1–20
- Hurteau M D, North M P, Koch G W and Hungate B A 2019 Opinion: managing for disturbance stabilizes forest carbon *Proc. Natl Acad. Sci.* **116** 10193
- Ibáñez I, Acharya K, Juno E, Karounos C, Lee B R, McCollum C, Schaffer-Morrison S and Tourville J 2019 Forest resilience under global environmental change: do we have the information we need? A systematic review *PLoS One* **14** e0222207
- Keith H, Mackey B, Berry S, Lindenmayer D and Gibbons P 2010 Estimating carbon carrying capacity in natural forest ecosystems across heterogeneous landscapes: addressing sources of error *Glob. Change Biol.* **16** 2971–89
- Körner C, Asshoff R, Bignucolo O, Hattenschwiler S, Keel S G, Pelaez-Riedl S, Pepin S, Siegwolf R T W and Zotz G 2005 Carbon flux and growth in mature deciduous forest trees exposed to elevated CO₂ *Science* **309** 1360–2
- Krug J H A 2019 How can forest management increase biomass accumulation and CO₂ sequestration? A case study on beech forests in Hesse, Germany *Carbon Balance Manage.* **14** 17
- Kurz W A *et al* 2009 CBM-CFS3: a model of carbon-dynamics in forestry and land-use change implementing IPCC standards *Ecol. Modelling* **220** 480–504
- Kurz W A, Shaw C H, Boisvenue C, Stinson G, Metsaranta J, Leckie D, Dyk A, Smyth C and Neilson E T 2013 Carbon in Canada's boreal forest—a synthesis *Environ. Rev.* **21** 260–92
- Lamb R L *et al* 2021 Context and future directions for integrating forest carbon into sub-national climate mitigation planning in the RGGI region of the U.S. *Environ. Res. Lett.* **16** 063001
- Lewis S L, Edwards D P and Galbraith D 2015 Increasing human dominance of tropical forests *Science* **349** 827–32
- Liang S, Hurteau M D and Westerling A L 2017 Potential decline in carbon carrying capacity under projected climate-wildfire interactions in the Sierra Nevada *Sci. Rep.* **7** 2420
- Liu J, Sleeter B M, Zhu Z, Heath L S, Tan Z, Wilson T S, Sherba J and Zhou D 2016 Estimating carbon sequestration in the piedmont ecoregion of the United States from 1971 to 2010 *Carbon Balance Manage.* **11** 1–13
- Markusson N 2022 Natural carbon removal as technology *WIREs Clim. Change* **13** e767
- Matthews H D *et al* 2020 Opportunities and challenges in using remaining carbon budgets to guide climate policy *Nat. Geosci.* **13** 769–79
- McIntire E J B, Chubaty A M, Cumming S G, Andison D, Barros C, Boisvenue C, Haché S, Luo Y, Micheletti T and Stewart F E C 2022 PERFICT: a re-imagined foundation for predictive ecology *Ecol. Lett.* **25** 1345–51
- McIntire E J B, Chubaty A, Luo Y and Cumming S G 2019 Develop and run spatially explicit discrete event simulation models 2.07 *R language* (available at: <https://spades.predictiveecology.org>, <https://github.com/PredictiveEcology/SpaDES>)
- Merganičová K *et al* 2019 Forest carbon allocation modelling under climate change *Tree Physiol.* **39** 1937–60
- Metsaranta J M, Shaw C, Kurz W A, Boisvenue C and Morken S 2017 Uncertainty of inventory-based estimates of the carbon dynamics of Canada's managed forest (1990–2014) *Can. J. For. Res.* **47** 1082–94
- Micheletti T *et al* 2021 Assessing pathways of climate change effects in SpaDES: an application to boreal landbirds of Northwest Territories Canada *Front. Ecol. Evol.* **9** 654
- Oberpriller J, Cameron D R, Dietze M C and Hartig F 2021 Towards robust statistical inference for complex computer models *Ecol. Lett.* **24** 1251–61
- Pappas C *et al* 2020 Aboveground tree growth is a minor and decoupled fraction of boreal forest carbon input *Agric. For. Meteorol.* **290** 108030
- Paradis G and LeBel L 2018 Estimating the value-creation potential of wood supply using a hybrid simulation-optimization approach *Technical Report CIRRELT-2018-24* (available at: www.cirrelt.ca/DocumentsTravail/CIRRELT-2018-24.pdf)
- Pilli R, Grassi G, Kurz W A, Viñas R A and Guerrero N H 2016 Modelling forest carbon stock changes as affected by harvest and natural disturbances. I. Comparison with countries' estimates for forest management *Carbon Balance Manage.* **11** 1–18
- Pilli R, Vizzarri M and Chirici G 2021 Combined effects of natural disturbances and management on forest carbon sequestration: the case of Vaia storm in Italy *Ann. For. Sci.* **78** 46
- Prescott C E, Katzensteiner K and Weston C 2021 Soils and restoration of forested landscapes *Soils and Landscape Restoration* ed J A Stanturf and M A Callaham (New York: Academic) ch 11, pp 299–331
- Running Steven W 2012 A measurable planetary boundary for the biosphere *Science* **337** 1458–9
- Sharma T, Kurz W A, Stinson G, Pellatt M G and Li Q 2013 A 100-year conservation experiment: impacts on forest carbon stocks and fluxes *For. Ecol. Manage.* **310** 242–55
- Simard S W, Roach W J, Defrenne C E, Pickles B J, Snyder E N, Robinson A and Lavkulich L M 2020 Harvest intensity effects on carbon stocks and biodiversity are dependent on regional climate in Douglas-fir forests of British Columbia *Front. For. Glob. Change* **3** 88
- Sleeter B M, Frid L, Rayfield B, Daniel C, Zhu Z and Marvin D C 2022 Operational assessment tool for forest carbon dynamics for the United States: a new spatially explicit approach linking the LUCAS and CBM-CFS3 models *Carbon Balance Manage.* **17** 1
- Smyth C E, Xu Z, Lemprière T C and Kurz W A 2020 Climate change mitigation in British Columbia's forest sector: GHG reductions, costs, and environmental impacts *Carbon Balance Manage.* **15** 21

- Sothe C, Gonsamo A, Arabian J and Snider J 2022 Large scale mapping of soil organic carbon concentration with 3D machine learning and satellite observations *Geoderma* **405** 115402
- Stall S, Yarmey L, Cutcher-Gershenfeld J, Hanson B, Lehnert K, Nosek B, Parsons M, Robinson E and Wyborn L 2019 Make scientific data FAIR *Nature* **570** 27–29
- Stinson G *et al* 2011 An inventory-based analysis of Canada's managed forest carbon dynamics, 1990–2008 *Glob. Change Biol.* **17** 2227–44
- Trouillier M, van der Maaten-theunissen M, Scharnweber T and Wilmking M 2020 A unifying concept for growth trends of trees and forests—the “potential natural forest” *Front. For. Glob. Change* **3** 113
- Wagner C E V 1978 Age-class distribution and the forest fire cycle *Can. J. For. Res.* **8** 220–7
- Walker W S *et al* 2022 The global potential for increased storage of carbon on land *Proc. Natl Acad. Sci.* **119** e2111312119
- Walker X J *et al* 2019 Increasing wildfires threaten historic carbon sink of boreal forest soils *Nature* **572** 520–3
- Wei S *et al* 2014 Data-based perfect-deficit approach to understanding climate extremes and forest carbon assimilation capacity *Environ. Res. Lett.* **9** 065002
- Xia J *et al* 2017 Terrestrial ecosystem model performance in simulating productivity and its vulnerability to climate change in the northern permafrost region *J. Geophys. Res.* **122** 430–46
- Zhao J, Daigneault A and Weiskittel A 2022 Estimating regional timber supply and forest carbon sequestration under shared socioeconomic pathways: a case study of Maine, USA *PLOS Clim.* **1** e0000018